

Table 22.10
Cancer of the Pancreas (Invasive)
SEER^a Relative Survival (Percent)
By Year of Diagnosis

All Races, Males

	Year of Diagnosis																			
	1975– 1979	1980– 1984	1985– 1989	1990– 1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
Survival Time																				
1-year	14.2	15.1	16.8	17.9	16.8	20.0	21.3	19.4	22.7	22.0	23.3	24.9	24.0	25.2	25.9	27.3	28.3	31.4	29.0	29.7
2-year	5.8	5.9	6.2	8.4	7.2	10.9	9.9	9.2	9.8	11.4	12.1	11.1	11.3	12.5	11.8	13.6	13.7	15.9	15.9	15.9
3-year	3.4	3.8	4.4	5.8	4.0	7.1	7.0	5.6	5.9	7.1	7.9	8.1	7.2	8.1	7.7	9.7	9.1	11.2		
4-year	2.9	3.0	3.2	4.6	3.4	5.8	5.5	4.6	5.5	5.9	6.3	6.8	5.3	5.7	5.5	8.1	7.8			
5-year	2.5	2.5	2.9	4.0	2.9	5.1	4.7	4.0	4.9	5.2	5.4	6.0	4.2	4.9	5.0	7.4				
6-year	2.4	2.2	2.4	3.8	2.6	4.6	3.9	3.8	4.0	4.9	5.0	5.2	3.8	4.3	4.6					
7-year	2.3	1.9	2.3	3.4	2.6	4.2	3.4	3.5	3.9	4.5	4.5	4.7	3.5	3.8						
8-year	2.0	1.7	2.1	3.1	2.4	4.2	3.4	3.3	3.9	3.9	4.1	4.3	3.4							
9-year	1.9	1.6	1.8	3.0	2.2	4.0	3.4	2.9	3.6	3.7	4.0	4.2								
10-year	1.8	1.5	1.7	2.7	2.1	3.8	3.2	2.9	3.6	3.5	3.8									
11-year	1.7	1.5	1.6	2.6	2.0	3.2	2.9	2.9	3.2	3.5										
12-year	1.6	1.5	1.6	2.6	1.8	3.0	2.8	2.6	3.1											
13-year	1.6	1.4	1.5	2.5	1.7	3.0	2.6	2.6												
14-year	1.6	1.4	1.5	2.3	1.2	2.9	2.4													
15-year	1.6	1.3	1.3	2.3	1.2	2.8														
16-year	1.6	1.3	1.3	2.1	1.2															
17-year	1.4	1.3	1.3	2.0																
18-year	1.2	1.3	1.2																	
19-year	1.1	1.3	1.2																	
20-year	1.1	1.2	1.2																	

^a Based on the SEER 9 areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, and Atlanta).